

# Ornithological Observations



An electronic journal published by BirdLife South Africa and the Animal Demography Unit at the University of Cape Town



Ornithological Observations accepts papers containing faunistic information about birds. This includes descriptions of distribution, behaviour, breeding, foraging, food, movement, measurements, habitat and plumage. It will also consider for publication a variety of other interesting or relevant ornithological material: reports of projects and conferences, annotated checklists for a site or region, specialist bibliographies, and any other interesting or relevant material.

**Editor: Arnold van der Westhuizen**

---

## **EVIDENCE OF BREEDING BY DIVING PETRELS AND STORM PETRELS AT MARION ISLAND AFTER THE ERADICATION OF FERAL CATS**

**Gregory TW McClelland, John Cooper and Stephen L Chown**

Recommended citation format:

**McClelland GTW, Cooper J, Chown SL 2013.** Evidence of breeding by Diving Petrels and Storm Petrels at Marion Island of the eradication of feral cats. *Ornithological Observations* Vol 4: 90-93

URL: <http://oo.adu.org.za/content.php?id=89>

Published online: 12 June 2013

**- ISSN 2219-0341 -**



## EVIDENCE OF BREEDING BY DIVING PETRELS AND STORM PETRELS AT MARION ISLAND AFTER THE ERADICATION OF FERAL CATS

Gregory TW McClelland, John Cooper\* and Steven L Chown

DST-NRF Centre of Excellence for Invasion Biology,  
Department of Botany and Zoology, Stellenbosch University, Private  
Bag X1, Matieland 7602, South Africa

\*Corresponding author: [john.cooper61@gmail.com](mailto:john.cooper61@gmail.com)

The South African Prince Edward Islands, comprising Marion Island (S46° 54', E37° 45') and the 21-km distant Prince Edward Island (S46° 38', E37° 57'), are among the most important sanctuaries for avifauna in the Southern Ocean. Twenty-nine species of bird have been recorded or are suspected to breed within the archipelago (Ryan and Bester 2008). Whereas the avifaunal community on near-pristine Prince Edward Island is considered to be essentially unaltered – it has never supported introduced predators – (De Villiers and Cooper 2008), Marion Island is likely still recovering from significant declines in its burrowing petrel populations as a consequence of four decades of predation by feral cats *Felis catus*.

Originally introduced to the Marion Island meteorological station in 1948 to control invasive House Mice *Mus musculus*, cats were first observed to have become feral and to feed on burrowing petrels in 1951/52 by RW Rand, the first biologist to make detailed observations of the island's birds (Rand 1954, Cooper 2008). They soon became well established around the entire island, having a deleterious effect on the island's bird populations. By 1977 cats were estimated to be taking approximately 455 000 burrowing petrels a

year (Van Aarde 1980). It was not until 1991 that the last cat was killed after an eradication effort spanning 14 years (Bester *et al.* 2000, 2002). Improved breeding success was subsequently reported for some of the larger burrowing petrels (Cooper and Fourie 1991, Cooper *et al.* 1995) but much less is known about the fortunes of the smaller burrowing species.

Rand (1954) described Common Diving Petrels *Pelecanoides urinatrix* (first reported breeding on Marion Island in 1948; Crawford 1952) as "...widely distributed over the coastal plain where they burrow under tussock and moss near the cliff edge" in 1951/52. However, active burrows could not be located during the 1965/66 summer despite "extensive" searches in the same sites by Van Zinderen Bakker Jr (1971) or subsequently in 1979/80 by Schramm (1986), suggesting that they had been eradicated as a breeding species by feral cats at an early stage. South Georgia Diving Petrels *P. georgicus*, which prefer to breed at higher altitudes on Marion Island where they were "commonly found" (Rand 1954), were still being recorded ("Many nests...found" and "eleven...nests opened one day...containing birds incubating eggs") in 1965/66 by Van Zinderen Bakker Jr (1971). Although the habitat preferred by this species (described as cinder slopes) was not well surveyed during a 1979/80 burrowing petrel survey of the eastern portion of the island, "very few burrows were found" (Schramm 1986), suggesting that this species was also succumbing to predation by cats. These few nests (estimated as only 44 burrows in 21 ha) were the last known record of a diving petrel breeding on Marion Island for the next 32 years.

On 10 March 2011 GTWM found an active diving petrel burrow at S46° 51.736', E37° 42.839' near the cinder cone Tumor at an elevation of 552 m. The burrow entrance was located under an *Azorella selago* cushion among unvegetated scoria and contained a




Fig 1- Diving petrel chick on Marion Island, 10 March 2011.

partially downy chick weighing 70 g. Although photographed (Fig .1), the chick was not definitively identified to species, although the altitude and substratum suggests it was a South Georgia Diving Petrel.

On 16 March 2009 two juvenile Black-bellied Storm-Petrels *Fregetta tropica*, a species yet to be confirmed as breeding on Marion (Cooper and Brown 1990), were captured on the same night by GTWM near Fur Seal Peninsula on the western side of the island. Conditions were extremely foggy at the time and the birds collided with GTWM as a result of being drawn in by a head torch. Both birds came from the direction of the interior and had traces of down on their abdomens. It is most likely that they had very recently fledged from the island and that the species breeds there, although it is conceivable they could have come from Prince Edward Island, where breeding has been confirmed (Berruti *et al.* 1981, Imber 1983). Black-bellied Storm-Petrels with "bare vascularized brood patches and enlarged gonads" have been caught on Marion Island in the past (Williams and Burger 1978) and adults have been caught occasionally by spot-lighting at night in the vicinity of the meteorological station in recent years (Oceans & Coasts, Department of Environmental Affairs unpubl. records).

The Grey-backed Storm-Petrel *Garrodia nereis* has never been definitely reported breeding on Marion Island, although it has been on Prince Edward Island (Van Zinderen Bakker Jr 1971, Cooper and Brooke 1984). The species has also been caught ashore on Marion with vascularized brood patches and enlarged gonads (Williams and Burger 1978, see also Klages *et al.* 1995).

- oo0oo -




### Acknowledgements

We thank the South African National Antarctic Programme, Department of Environmental Affairs for logistic support and the South African National Research Foundation for financial support via grant SNA2007044200003 to SLC. We thank RJM Crawford of the Department of Environmental Affairs for providing unpublished bird records. PG Ryan provided valuable comments that improved the manuscript.

### References

- Berruti A, Griffiths AM, Imber MJ, Schramm M, Sinclair JC 1981.** The status of seabirds at Prince Edward Island. South African Journal of Antarctic Research 10/11: 31-32.
- Bester MN, Bloomer JP, Bartlett PA, Muller DD, Van Rooyen M, Büchner H 2000.** Final eradication of feral cats from sub-Antarctic Marion Island, southern Indian Ocean. South African Journal of Wildlife Research 30: 53-57.
- Bester MN, Bloomer JP, Van Aarde RJ, Erasmus BH, Van Rensburg PJJ, Skinner JD, Howell PG, Naude TW 2002.** A review of the successful eradication of feral cats from sub-Antarctic Marion island, southern Indian Ocean. South African Journal of Wildlife Research 31: 65-73.
- Cooper J 2008.** Human history. In: Chown SL, Froneman PW (eds). The Prince Edward Islands. Land-sea interactions in a changing ecosystem. SUN PReSS, Stellenbosch: 331-350.
- Cooper J, Brooke MdeL 1984.** Breeding status of burrowing petrels at Prince Edward Island. South African Journal of Antarctic Research 14: 34-35.
- Cooper J, Brown CR 1990.** Ornithological research at the sub-Antarctic Prince Edward Islands: a review of achievements. South African Journal of Antarctic Research 20: 40-57.
- Cooper J, Fourie A 1991.** Improved breeding success of Great-winged Petrels *Pterodroma macroptera* following control of feral cats *Felis catus* at subantarctic Marion Island. Bird Conservation International 1: 171-175.
- Cooper J, Marais AVN, Bloomer JP, Bester MN 1995.** A success story: breeding of burrowing petrels (Procellariidae) before and after the eradication of feral cats *Felis catus* at subantarctic Marion Island. Marine Ornithology 23: 33-37.
- Crawford AB 1952.** The birds of Marion Island, South Indian Ocean. Emu 52: 73-85.
- De Villiers MS, Cooper J 2008.** Conservation and management. In: Chown SL, Froneman PW (eds). The Prince Edward Islands. Land-sea interactions in a changing ecosystem. SUN PReSS, Stellenbosch: 301-330.
- Imber MJ 1983.** The lesser petrels of the Antipodes Islands, with notes from Prince Edward and Gough Islands. Notornis 30: 283-298.
- Klages NTW, Nel DC, Cooper J 1995.** Stomach contents of a Grey-backed Storm Petrel *Garrodia nereis* from Subantarctic Marion Island. Marine Ornithology 23: 163-164.
- Rand RW 1954.** Notes on the birds of Marion Island. Ibis 96: 173-206.
- Ryan PG, Bester MN 2008.** Pelagic predators. In: Chown SL, Froneman PW (eds). The Prince Edward Islands. Land-sea interactions in a changing ecosystem. SUN PReSS, Stellenbosch: 121-164.



---

**Schramm M 1986.** Burrow densities and nest site preferences of petrels (Procellariidae) at the Prince Edward Islands. *Polar Biology* 6: 63-70.

**Van Aarde RJ 1980.** The diet and feeding behaviour of feral cats, *Felis catus* at Marion Island. *South African Journal of Wildlife Research* 10: 123-128.

**Van Zinderen Bakker Jr EM 1971.** Comparative avian ecology. In: Van Zinderen Bakker Sr EM, Winterbottom JM, Dyer RA (eds). Marion and Prince Edward Islands. Report on the South African Biological & Geological Expedition /1965-1966. A.A. Balkema, Cape Town: 161-172.

**Williams AJ, Burger AE 1978.** Notes on nonbreeding seabirds at the Prince Edward Islands. *Cormorant* 5: 11-14.